


PLANT GROWTH & DEVELOPMENT

- All plants $\xrightarrow[\text{of}]{\text{descendants}}$ Zygote
- Growth + Differentiation = Development (sum)
- Zygote $\xrightarrow{\text{follow a precise \& highly ordered succession of events}}$ Mature plant (complex body organ. - Roots, Leaves, branch, flower)



GROWTH : \rightarrow conspicuously involves increased protoplasmic material.

- Most fundamental & conspicuous charact. of living beings.
- ① Irreversible ② permanent increase in size of an organ or its parts or even of an indiv. cell.

- Growth $\xrightarrow{\text{accompanied by}}$ Metabolic activities \rightarrow occur at the expense of energy.
 - \swarrow both \searrow
 - Catabolic Anabolic

Example \rightarrow Expansion of leaf.

- PLANT GROWTH GENERALLY IS INDETERMINATE : \rightarrow Imp.

Plant growth $\xrightarrow{\text{Unique}}$ Retain the capacity for unlimited growth throughout life

because of .
 presence of meristems at certain locations \rightarrow capacity to divide & self perpetuate in their body.

Root apical meristem
Shoot apical meristem
Intercalary meristem

contribute to elongation
growth of plant ^{ones}.


Date
Page

The product → loses the capacity to divide
& such cells make up plant body.

* open form of growth present

→ Root & shoot apical meristem. → primary growth of the plants → elongation along plant axis.

→ Dicots & Gymnosperm → Lateral meristems



TANISHA SACHAN

AIR 1747

GROWTH IS MEASURABLE

NCERT THREAD NOTES

Growth → at cellular level → consequence → increase in amt of protoplasm

- 1) increase in
① fresh weight
② dry weight
③ Length ④ Area
⑤ Volume ⑥ Cell number.

difficult to measure directly
hence one measures some
quantity which is more
or less proportional to it.

→ One single Maize Root apical meristem → give rise → > 17,500 new cells per hour
Increase in cell number.

→ Cells of watermelon → may increase in size → upto 3,50,000 times
Increase in size of cell

→ Growth of pollen tube → In terms of length

→ Growth in adorsiventral, f → in terms of surface area.

Phases of Growth:

Meristematic

- Cells rich in protoplasm
- Relatively large conspicuous nuclei
- Cell walls - primary thin cellulose
With plasmodesmata connection (many)

Eg. Constantly dividing cells, both at root apex & shoot apex.

Elongation

- Increased vacuolation
- Cell enlargement
- New cell wall deposition

Eg. Cells proximal to meristematic zone.

Maturation

- Cells in maximal size in terms of ↓
- Wall thickening
- Protoplasmic modifications

Eg. Cells proximal to elongation zone

Proximal
→ away from tip.

Growth Rates:

Growth ↑ per unit time

Two types of growth:

ARITHMETIC

Following mitotic cell division only one daughter cell divide and other differentiates & matures.

Eg. Root elongating at a constant rate.

$$L_t = L_0 + \gamma t$$

growth rate/
elong. per unit time

GEOMETRIC

In initial stage, growth is slow (Lag phase)

then ↑ rapidly at exponential rate (log/exponential phase)

Here both the progeny cells follow mitotic cell division & retain the ability to divide & so on.

With limited nutrient supply, growth slows down (Stationary phase)

$$W_t = W_0 e^{\gamma t}$$

efficiency index

Relative growth rate/measure of the ability of plant to prod. new plant material
base of natural logarithms


DEVELOPMENT:

↓ Includes


NCERT THREAD NOTES

All changes that org. goes throughout life cycle


* Applicable to tissues & organ also.



Plants follow diff. pathway → In response to environ. or phases of life → To form different kind of structures (This ability is plasticity)



* Thus Growth, differentiation & development very closely related events in plant life.





Roles of PGR could be Complimentary Or antagonistic.
These could be individualistic or synergistic.

Role of PGR → intrinsic control

→ Along with genomic & external factors

Imp role in plant growth & development.

Many intrinsic factors such as temp & light control plant growth & develop via PGR

1) Vernalisation

2) Flowering

3) Dormancy

4) Seed germination

5) Plant movement.

PHOTO PERIODISM :

→ The critical duration is diff. for differ. plants.

- Some plants require periodic exposure to light to induce flowering.
- Plants are able to measure duration of light

Long day plants

day-neutral plant

Short day plants

* Duration of dark period is also important.

* Flowering in certain plants depends not only on the combination of light & dark exposures but also their relative durations.


- The response of plants to periods of day/night.



Date

Page

→ shoot apices ^{modify} → flowering apices
(before flowering)
→ cannot perceive photoperiods

★ The site of perception of light/dark duration are →  Leaves.

Hypothesized → hormonal substance → responsible for flowering
→ migrate from Leaves → shoot apices

TANISHA SACHAN

AIR 1747

NCERT THREAD NOTES

VERNALISATION → Promotion of flowering by a period of low temperature

Flowering - Quantitatively or Qualitatively dependent on exposure to low temp

* It prevents precocious reproductive development late in the growing season, and, enables the plant to have sufficient time to reach maturity.

Imp Food plants: "Wheat & Barley"

"Rye"

have two kinds of varieties

Winter variety

Spring variety

If planted in spring would normally fail to flower / or produce mature grain within a span of flowering season.

Planted in: spring
Flower/grain in: Before the end of growing season.

Planted in: Autumn.

Germinate: over winter

* Come out as small seedlings

Resume growth: in Spring

Harvested: Around mid summer.

Another example of vernalisation

Biennial plants

Sugarbeet
Cabbages
Carrots

Are MONOCARPIC

Normally flower & die in 2nd season.

* Subjecting the growing of biennial plant to a cold treatment stimulates a subsequent photoperiodic flowering response.

TANISHA SACHAN

SEED DORMANCY

AIR 1747

NCERT THREAD NOTES

Certain seeds → fail to germinate → when external cond. are favourable.

Such seeds are going a period of dormancy

Controlled 'not' by external env. but are "under" endogenous control or cond. within seeds itself.

Reasons which causes dormancy

① Impermeable & hard seed coat.


② Presence of chemical inhibitors such as
→ abscisic acid
→ phenolic acid
→ Para-ascorbic acid.

③ Immature embryos.

* Plants exhibit plasticity in development.

PGR - control different differentiation & development events.
 → have diverse physiological effect on plant.

Overcoming Dormancy



(*) Effect of inhibitory substance can be removed by subjecting the seeds to

- chilling cond.
- Application of chemicals
 - GA
 - Nitrate

(*) Changing Environment cond such as

- Light
- Temp.


→ Other ways to overcome seed dormancy

(*) Growth may not be & generally is not sustained at high rate through the life of cell/tissue/organ/organism.

(*) Differentiation results in development of structures, that is commensurate with the function the cells finally has to perform.

General principles for differentiation of cells, tissue & organs are similar.


small, simple molecule
of diverse chemical composition.



TANISHA SACHAN


AIR 1747

NCERT THREAD NOTES



Discovery of PGR.

"Discovery of each of the five PGR. was have been accidental."



Abscicic acid

In mid 1960s

Three independ. researchers

↓ reported

purification & chemical
characterisation
of three diff. kinds of
inhibitors

↓
inhibitor-B dormin abscission II

Later all three were proved
chemically identical

↓
ABA (abscicic acid)

Ethylene

H. H. Cousins (1910)

confirmed release of
volatile subset

↓

from ripened oranges.

that hastened ripening of
unripe bananas.

↓
Ethylene
(gaseous)

AUXIN (auxein: to grow) by greek.

Date _____
Page _____


TANISHA SACHAN

AIR 1747

NCERT THREAD NOTES

→ Isolated from : human urine

→



Higher plants → Growing apical buds inhibit growth of lateral bud.
(Apical dominance)

Removal of shoot tips
(decapitation)

Growth of lateral buds.

Widely applied in tea plantations, hedge making.

Natural

- IAA
- IBA

Isolated from plants

Synthetic

• NAA
(Naphthalene acetic acid)

• 2,4-D
(2,4-dichlorophenoxyacetic)

* All these auxins have been used extensively in 1) agricultural 2) Horticulture.

→ Auxin → Initiate rooting in stem cuttings
→ Plant propagation.

→ Flowering : In pineapples

→ Prevent fruit & leaf drop at early stage.

→ Promote abscissions of older mature leaves & fruits

→ Parthenocarpy : In tomatoes.

→ Herbicides

↓
2,4-D

Widely used to kill dicotyledonous weeds plants but does not affect dicot plants.

↓
Used to prepare weed free lawns by gardeners.

Xylem differentiation


Cell division

GIBBERELLINS

TANISHA SACHAN
AIR 1747

NCERT THREAD NOTES

- Promotory PGR.



* GAs produce a variety of biological response in plants.

- ability to increase in length of axis
 - To increase length of grape stalks.
- Apple, to elongate & improve its shape.
- Delays senescence. (thus fruits can be left on tree longer to extend market period)
- Speed up malting process in brewing industry.

Sugarcane : stores carbohydrates (as sugars) in their stems.

Spraying with gibberellins → ↑ length of stem.

↑ yield by 20 tonnes per acre.

Juvenile Conifers : Spraying these with gibberellins



Hastens maturity period → thus → Early (speedy) seed production.

Beet, Cabbage & Plants with Rosette habit

: GA promotes bolting (internode elongations just prior to flowering)

CYTOKININ

Richmond Lang Effect.

- Have special effects on cytokinesis
- Was discovered as Kinetin (a modified form of adenine, purine)

TANISHA SACHAN

AIR 1747

NCERT THREAD NOTES

does not occur naturally in plants.

Autoclaved herring sperm DNA.

↑ (from)

Search for subst. (natural) with cytokinin like activities

led to

Isolation of zeatin

from

① Corn kernel ② Coconut milk.

- Since zeatin's discovery, several natural as well as synthetic comp. with cell division promoting activity have been identified

- Natural cytokinin found out in

region where rapid cell division occurs.

- Root apices
- Developing shoot buds
- Developing/young fruits

- Cytokinin helps to produce → new leaves

new chloroplast in leaves

Lateral shoot growth

adventitious shoot formation.

- Cytokinins help overcome apical dominance.


- Help in nutrient mobilisation — helps in — delay of leaf senescence.

ETHYLENE

NCERT THREAD NOTES

- Simple, gaseous PGR.
- Synthesized (in large quantities) by : Tissue undergoing senescence
Ripening fruits.

- Its influences →
 - Horizontal growth of seedlings
 - Swelling of the axis
 - Apical hook formation in dicot seedlings



- Source : Ethephon → aq. soln → readily absorbed & transported within the plant & releases ethylene slowly.

- Ethylene hastens fruit ripening
Tomato Apples

① accelerates abscission in flowers & fruit (thinning of cotton, cherry, walnut)

① Promotes female flowers in cucumbers & thereby increasing yield

ABSCISIC ACID




Date / /
Page

- Has role in regulating abscission & dormancy.
- Plant growth inhibitor & an inhibitor of plant metabolism.
- Inhibits seeds germination.
- Stimulates closure of stomata in epidermis

↓
Increases tolerance of plants to various kind of stress

↓
"Stress hormone"



TANISHA SACHAN

AIR 1747

NCERT THREAD NOTES

* Both Long day & Short day plant - can produce flower simultaneously in given place

* Cytokinin - cz senescence of leaf